

# Publisher's Note: "The Landau-de Gennes approach revisited: A minimal self-consistent microscopic theory for spatially inhomogeneous nematic liquid crystals" [J. Chem. Phys. 147, 244505 (2017)]

Cite as: J. Chem. Phys. 148, 029901 (2018); <https://doi.org/10.1063/1.5021484>  
Submitted: 29 December 2017 • Published Online: 10 January 2018

 Ioana C. Gârlea and  Bela M. Mulder



View Online



Export Citation



CrossMark

## ARTICLES YOU MAY BE INTERESTED IN

[The Landau-de Gennes approach revisited: A minimal self-consistent microscopic theory for spatially inhomogeneous nematic liquid crystals](#)

The Journal of Chemical Physics **147**, 244505 (2017); <https://doi.org/10.1063/1.4993574>

[Defects in liquid crystals](#)

Physics Today **35**, 48 (1982); <https://doi.org/10.1063/1.2915094>

[A novel model for smectic liquid crystals: Elastic anisotropy and response to a steady-state flow](#)

The Journal of Chemical Physics **145**, 164903 (2016); <https://doi.org/10.1063/1.4965711>

 **The Journal of Chemical Physics** **Special Topics** Open for Submissions [Learn More](#)

## Publisher's Note: "The Landau-de Gennes approach revisited: A minimal self-consistent microscopic theory for spatially inhomogeneous nematic liquid crystals" [J. Chem. Phys. **147**, 244505 (2017)]

Ioana C. Gârlea<sup>a)</sup> and Bela M. Mulder

*Institute AMOLF, Science Park 104, 1098XG Amsterdam, The Netherlands*

(Received 29 December 2017; published online 10 January 2018)

<https://doi.org/10.1063/1.5021484>

This article was originally published online on 26 December 2017 with three errors in Eq. (1). Three centered dots should have appeared as wedge symbols. AIP Publishing apologizes for this error. All online versions of the article were corrected on 3 January 2018; the article is correct as it appears in the printed version of the journal.

The correct Eq. (1) appears below:

$$E[\hat{\mathbf{n}}] = \frac{1}{2} \int d\mathbf{r} \left\{ K_1 (\nabla \cdot \hat{\mathbf{n}}(\mathbf{r}))^2 + K_2 (\hat{\mathbf{n}}(\mathbf{r}) \cdot \nabla \wedge \hat{\mathbf{n}}(\mathbf{r}))^2 + K_3 (\hat{\mathbf{n}}(\mathbf{r}) \wedge \nabla \wedge \hat{\mathbf{n}}(\mathbf{r}))^2 \right\}. \quad (1)$$

---

<sup>a)</sup>Current address: Faculty of Physics, University of Vienna, Boltzmannngasse 5, A-1090 Vienna, Austria.